

# Test Report: BS EN 14476:2013 + A2:2019 Chemical disinfectants and antiseptics – Quantitative suspension test for the evaluation of virucidal activity in the medical area- Test method and requirements (Phase 2/Step 1)

**Test Laboratory****BluTest Laboratories Ltd**

5 Robroyston Oval, Nova Business Park, Glasgow, G33 1AP

**Identification of sample**

|                        |   |
|------------------------|---|
| Name of the product    | <b>F10SC Veterinary Disinfectant</b>          |
| Batch number           | 220309  |
| Client                 | Health and Hygiene (Pty) Limited              |
| Client Address         | PO Box 906, Florida Hills, 1716, South Africa |
| Project Code           | BT-HAH-02                                     |
| Date of Delivery       | 02 July 2020                                  |
| Storage conditions     | Ambient                                       |
| Active substances      | QAC and PHMB 5.8%                             |
| Appearance             | Liquid  |
| Condition upon receipt | Undamaged                                     |

**Test Method and its validation**

|                |   |
|----------------|---|
| Method         | 1 part interfering substance + 1 part virus suspension + 8 parts biocide were mixed and incubated at the indicated contact temperature for the indicated contact times. Assays were validated by a cytotoxicity control, interference control, neutralisation control and a formaldehyde internal standard. |
| Neutralisation | Dilution-neutralisation/gel filtration<br>Eagles Minimum Essential Medium + 5.0% v/v foetal bovine serum at 4°C   |

**Experimental Conditions**

|   |   |
|---|---|
| Period of analysis                      | 11 August 2020 to 16 August 2020                  |
| Product diluents used                   | Sterile, synthetic hard water                     |
| Product test concentrations             | 1:1500; 1:500; 1:250                              |
| Appearance product dilutions            | No changes noted- stable                          |
| Appearance in test mixture              | Sedimentation and turbidity observed at 1:250     |
| Contact times (minutes)                 | 5 ± 10s   |
| Test temperature                        | 20°C ± 1°C  |
| Interfering substances                  | 0.3g/l bovine albumin                             |
| Temperature of incubation               | 37°C ± 1°C + 5% CO <sub>2</sub>                   |
| Identification and passage (P) of virus | <b>Vaccinia virus VR-1549 Elstree strain (P7)</b> |
| Identification and passage (P) of cells | Vero Cells (P 42)                                 |

## PROTOCOL SUMMARY

The basic virucidal efficacy test is set up with three concentrations of test product solution and a 5 minute contact time. Virus is exposed to disinfectant in 24-well plates, then neutralised, serially diluted and virus titred in 96-well tissue culture plates to determine the tissue culture infectious dose<sub>50</sub> (TCID<sub>50</sub>) of surviving virus. *Vaccinia virus* VR-1549 Elstree strain / Vero cells are assayed in parallel in each test. TCID<sub>50</sub> is determined by the method of Karber<sup>1</sup>.

### **Cytotoxicity control**

The test product solution is measured for its effects on the host cells used to propagate the virus, to determine the sensitivity of the assay.

### **Interference control**

The effect of the cells after treatment of the test product solution are verified to ensure the cells can show susceptibility for virus infection. This is compared against cells that have not been treated with test product.

### **Disinfectant suppression control VS1**

Virus is added to the highest concentration of test product solution and then the mixture immediately removed and neutralised. The neutralised virus titre is then determined to assess the efficiency of the neutralisation procedure.

### **Disinfectant suppression control VS2**

Internal control which adds virus to neutralised test product solution to assess the efficiency of the neutralisation procedure.

### **No column Control**

Internal control on the highest contact time to assess any impact of the Microspin™ S 400 HR columns.

### **Virus recovery control**

Virus titre is determined for virus in contact with sterile, synthetic hard water at t=0, t = 5 and at t =15. The virus titre after 5 minutes is then compared to the recovery of disinfectant-treated virus to measure the log reduction in virus titre. The virus titre at 15 minutes is compared to the reference virus inactivation control.

### **Reference virus inactivation control**

Virus is exposed to 0.7% W/V formaldehyde and the recovery of virus determined by TCID<sub>50</sub> after 5 and 15 minutes, in order to assess that the test virus has retained reproducible biocide resistance. In addition, the formaldehyde cytotoxicity of neutralised formaldehyde is determined, to measure assay sensitivity.

1Kärber, G.: Beitrag zur Kollektiven Behandlung Pharmakologischer Reihenversuche. Arch. Exp. Path. Pharmak. 162 (1931): 480-487.

### Vaccinia virus (VR-1549) Elstree strain Test Results

| EN14476:2013 + A2:2019 Suspension test for the efficacy of F10SC Veterinary Disinfectant, BT-HAH-02 from Health and Hygiene (Pty) Limited against Vaccinia virus ATCC VR-1549 under CLEAN conditions |        |                        |        |                        |        |                        |
|--|--------|------------------------|--------|------------------------|--------|------------------------|
| Test Results   |        |                        |        |                        |        |                        |
| Concentration  | 1:1500 |                        | 1:500  |                        | 1:250  |                        |
| Exposure Time  | data   | TCID <sub>50</sub> /ml | data   | TCID <sub>50</sub> /ml | data   | TCID <sub>50</sub> /ml |
| t = 5 minutes  | 4.17   | 4.64E+05               | 3.17   | 4.64E+04               | 0.00   | 3.16E+01               |
| Raw Data   | 666610 | 4.64E+05               | 666100 | 4.64E+04               | 000000 | 3.16E+01               |
| log  |        | 5.67                   |        | 4.67                   |        | 1.50                   |
| log difference   |        | 0.83                   |        | 1.83                   |        | 5.00                   |

| EN14476:2013 + A2:2019 Suspension test for the efficacy of F10SC Veterinary Disinfectant, BT-HAH-02 from Health and Hygiene (Pty) Limited against Vaccinia virus ATCC VR-1549 under CLEAN conditions |                       |               |                       |                       |       |        |        |        |                               |
|--|-----------------------|---------------|-----------------------|-----------------------|-------|--------|--------|--------|-------------------------------|
| Summary Table  |                       |               |                       |                       |       |        |        |        |                               |
| Product:   | Interfering substance | Concentration | Level of cytotoxicity | lg TCID <sub>50</sub> |       |        |        |        | >4 lg reduction after 'X' Min |
|  |                       |               |                       | 0 min                 | 5 min | 15 min | 30 min | 60 min |                               |
| F10SC Veterinary Disinfectant  | 0.3g/l BSA            | 1:250         | 1.50                  | 4.67                  | 1.50  | n.a.   | n.a.   | n.a.   | < 5 minutes                   |
|  |                       | 1:500         | 1.50                  | n.a.                  | 4.67  | n.a.   | n.a.   | n.a.   | > 5 minutes                   |
|  |                       | 1:1500        | 1.50                  | n.a.                  | 5.67  | n.a.   | n.a.   | n.a.   | > 5 minutes                   |
| Virus Control  | CLEAN                 |               |                       | 6.50                  | 6.50  | 6.50   | 6.50   | 6.50   | n.a.                          |
|  |                       |               |                       |                       |       |        | 5 min  | 15 min |                               |
| Formaldehyde   | PBS                   | 0.7% (w/v)    | 2.50                  |                       |       |        | 4.67   | 2.50   | >15 mins                      |

### Vaccinia virus (VR-1549) Elstree strain Control Data

EN14476:2013 + A2:2019 Suspension test for the efficacy of F10SC Veterinary Disinfectant, BT-HAH-02 from Health and Hygiene (Pty) Limited against Vaccinia virus ATCC VR-1549 under CLEAN conditions

| Controls                                     |                        |                      |                        |                        |                        |                        |                        |                                   |                        |                              |                        |  |
|--|------------------------|----------------------|------------------------|------------------------|------------------------|------------------------|------------------------|-----------------------------------|------------------------|------------------------------|------------------------|--|
| Virus Recovery 0 min                         |                        | Virus Recovery 5 min |                        | Virus Recovery 15 min  |                        | Cytotoxicity           |                        | Disinfectant Suppression VS       |                        | Disinfectant Suppression VS2 |                        |  |
| raw data                                     | TCID <sub>50</sub> /ml | raw data             | TCID <sub>50</sub> /ml | raw data               | TCID <sub>50</sub> /ml | raw data               | TCID <sub>50</sub> /ml | raw data                          | TCID <sub>50</sub> /ml | raw data                     | TCID <sub>50</sub> /ml |  |
| 5.00   | 3.16E+06               | 5.00                 | 3.16E+06               | 5.00                   | 3.16E+06               | 0.00                   | 3.16E+01               | 3.17                              | 4.64E+04               | 5.00                         | 3.16E+06               |  |
| 666660                                       | 3.16E+06               | 666660               | 3.16E+06               | 666651                 | 3.16E+06               | 000000                 | 3.16E+01               | 665200                            | 4.64E+04               | 666651                       | 3.16E+06               |  |
|  | 6.50                   |                      | 6.50                   |                        | 6.50                   |                        | 1.50                   |                                   | 4.67                   |                              | 6.50                   |  |
|  |                        |                      |                        |                        |                        |                        |                        |                                   | 1.83                   |                              | 0.00                   |  |
| Formaldehyde reference inactivation controls |                        |                      |                        |                        |                        |                        | No column Control      |                                   |                        |                              |                        |  |
| Cytotoxicity                                 |                        | Exposure time        | 0.7% Formaldehyde      |                        |                        |                        | 5 mins                 |                                   | 15 mins                |                              | 5 mins                 |  |
| raw data                                     | TCID <sub>50</sub> /ml |                      | raw data               | TCID <sub>50</sub> /ml | raw data               | TCID <sub>50</sub> /ml | raw data               | TCID <sub>50</sub> /ml            | raw data               | TCID <sub>50</sub> /ml       | TCID <sub>50</sub> /ml |  |
| 1.00   | 3.16E+02               |                      | 3.17                   | 4.64E+04               | 1.00                   | 3.16E+02               |                        |                                   | 5.33                   |                              | 6.81E+06               |  |
| 600000                                       | 3.16E+02               |                      | 666100                 | 4.64E+04               | 600000                 | 3.16E+02               |                        |                                   | 666662                 |                              | 6.81E+06               |  |
|  | 2.50                   | log                  |                        | 4.67                   |                        | 2.50                   |                        |                                   |                        |                              | 6.83                   |  |
|  |                        | log difference       |                        | 1.83                   |                        | 4.00                   |                        |                                   |                        |                              |                        |  |
| Interference control                         |                        | Virus dilution       |                        |                        |                        |                        |                        | Stock Virus (TCID <sub>50</sub> ) |                        |                              |                        |  |
|  |                        | -3                   | -4                     | -5                     | -6                     | -7                     | -8                     |                                   |                        |                              |                        |  |
| PBS Control                                  |                        | 1                    | 1                      | 1                      | 1                      | 0.33                   | 0                      | 6.33                              |                        |                              |                        |  |
|  |                        | 3.16E+02             | 3.16E+02               | 3.16E+02               | 3.16E+02               | 6.76E+01               | 3.16E+01               | 6.76E+07                          |                        |                              |                        |  |
|  |                        | 2.50                 | 2.50                   | 2.50                   | 2.50                   | 1.83                   | 1.50                   | 6666662000                        |                        |                              |                        |  |
| Raw Data                                     |                        | 6                    | 6                      | 6                      | 6                      | 2                      | 0                      |                                   |                        |                              |                        |  |
| Product                                      |                        | 1                    | 1                      | 1                      | 0.83                   | 0                      | 0                      |                                   |                        |                              |                        |  |
|  |                        | 3.16E+02             | 3.16E+02               | 3.16E+02               | 2.14E+02               | 3.16E+01               | 3.16E+01               |                                   |                        |                              |                        |  |
|  |                        | 2.50                 | 2.50                   | 2.50                   | 2.33                   | 1.50                   | 1.50                   |                                   |                        |                              |                        |  |
| Raw Data                                     |                        | 6                    | 6                      | 6                      | 5                      | 0                      | 0                      |                                   |                        |                              |                        |  |
| Log Difference                               |                        | 0.00                 | 0.00                   | 0.00                   | 0.17                   | 0.33                   | 0.00                   |                                   |                        |                              |                        |  |
| Product Cyt Dilution                         |                        | -1                   | -1                     | -1                     | -1                     | -1                     | -1                     |                                   |                        |                              |                        |  |
| PBS Dilution                                 |                        | Neat                 | Neat                   | Neat                   | Neat                   | Neat                   | Neat                   |                                   |                        |                              |                        |  |

## CONCLUSION

### Verification of the methodology

A test is only valid if the following criteria are fulfilled:

- a) The titre of the test suspension of at least  $10^8$  TCID<sub>50</sub> /ml is sufficiently high to at least enable a titre reduction of 4 lg to verify the method.
- b) Detectable titre reduction is at least 4 log<sub>10</sub>.
- c) Difference of the logarithmic titre of the virus control minus the logarithmic titre of the test virus in the reference inactivation test is between:
  - Between 0.75 and 3.5 after 5 min and between 2.0 and 4.0 after 15 min for Vaccinia virus
- d) Cytotoxicity of the product solution does not affect cell morphology and growth or susceptibility for the test virus in the dilutions of the test mixtures which are necessary to demonstrate a 4 log<sub>10</sub> reduction of the virus.
- e) The interference control result does not show a difference of > 1.0 log<sub>10</sub> of virus titre for test product treated cells in comparison to the non-treated cells.
- f) Neutralisation validation. This is called the disinfectant suppression test in this protocol. The disinfectant was neutralised by column chromatography through an Illustra Microspin S-400 HR column to achieve the best possible neutralisation available for this test. The difference for virus is greater than 0.5 log<sub>10</sub> indicating rapid irreversible virucidal activity of the disinfectant by dilution at a concentration of 1:250 for VS1. This neutralisation validation has been verified by VS2, which shows the product has been successfully neutralised.

According to EN 14476:2013 + A2:2019, **F10SC Veterinary Disinfectant POSSESSES VIRUCIDAL** activity at a concentration of **1:250** as tested after **5 MINUTES** at **20°C** under **CLEAN** conditions (0.3 g/l bovine albumin) against *Vaccinia virus* VR-1549 Elstree strain / Vero cells.

**This product therefore is effective against all enveloped viruses as defined in EN 14476:2013 + A2:2019 Annex A\*. This therefore includes all coronaviruses and SARS-CoV-2.**

Authorised signatory



Dr Chris Woodall, Director  
BluTest Laboratories Ltd  
Glasgow, UK  
Date: 03 September 2020

### DISCLAIMER

The results in this test report only pertain to the sample supplied.

BluTest (BT) has performed the testing detailed in this report using reasonable skill and care and has used reasonable endeavours to carry out the testing in accordance with an EN 14476 protocol. All forecasts, recommendations and results contained in this report are submitted in good faith. However, other than as expressly set out in this report, no warranty is given (i) in relation to the testing or the use(s) to which any results or deliverables produced in the course of the testing are or may be put by the Client or their fitness or suitability for any particular purpose or under any special conditions notwithstanding that any such purpose or conditions may have been made known to BT or (ii) that the intended results or deliverables from the testing can be achieved or (iii) that the Client can freely make use of the results or the deliverables without infringing any third party intellectual property rights and the Client will be deemed to have satisfied itself in this regard. BT shall have no liability (which is hereby excluded to the fullest extent permissible by law) in respect of any loss, liability or damage, including without limitation any indirect and/or consequential loss such as loss of profit or loss of business, market or goodwill, that the Client may suffer directly or indirectly as a result of or in connection with: (i) the performance of the testing; (ii) the use of any materials, samples or other information provided by the Client for use in the testing; and (iii) the Client's reliance upon or use of any results or deliverables provided as part of the testing.

**\*EN 14476 2013 + A2 2019 Annex A (informative – Enveloped viruses)**

Poxviridae  
Herpesviridae  
Filoviridae (e.g. Ebola, Marburg)  
Flavivirus  
Hepatitis C Virus (HCV)  
Hepatitis Delta Virus (HDV)  
Influenza Virus  
Paramyxoviridae  
Rubella Virus  
Measles Virus  
Rabies Virus  
Coronavirus (e.g. SARS, MERS)  
Human Immunodeficiency Virus (HIV)  
Human T Cell Leukemia Virus (HTLV)  
Hepatitis B virus (HBV)

Reference: Van Regenmortel MHV et al.,Eds.: Virus Taxonomy, Classification and Nomenclature of Viruses, seventh report of the international committee on taxonomy of viruses. Academic Press, San Diego, 2000